

Synonym

CD38,T10,cADPr hydrolase 1

Source

Human CD38 Protein, Llama IgG2b Fc Tag(CD8-H5252) is expressed from human 293 cells (HEK293). It contains AA Val 43 - Ile 300 (Accession # [P28907-1](#)).
Predicted N-terminus: Val 43

Molecular Characterization

CD38(Val 43 - Ile 300) P28907-1	LlamaFc(Glu1 - Ser243) AAX73259.1
------------------------------------	--------------------------------------

This protein carries a llama IgG2b Fc tag at the C-terminus.
The protein has a calculated MW of 57.7 kDa. The protein migrates as 66-90 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.01 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human CD38 Protein, Llama IgG2b Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA





Immobilized Human CD38 Protein, Llama IgG2b Fc Tag (Cat. No. CD8-H5252) at 1 µg/mL (100 µL/well) can bind Anti-CD38 MAb, Human IgG1 with a linear range of 0.4-6 ng/mL (QC tested).

Background

CD antigen CD38 is also known as ADP-ribosyl cyclase 1, which belongs to the ADP-ribosyl cyclase family. CD38 is expressed at high levels in pancreas, liver, kidney, brain, testis, ovary, placenta, malignant lymphoma and neuroblastoma. CD38 is a multifunctional ectoenzyme that catalyzes the synthesis and hydrolysis of cyclic ADP-ribose (cADPR) from NAD⁺ to ADP-ribose. These reaction products are essential for the regulation of intracellular Ca²⁺. The loss of CD38 function is associated with impaired immune responses, metabolic disturbances, and behavioral modifications. The CD38 protein is a marker of cell activation. It has been connected to HIV infection, leukemias, myelomas, solid tumors, type II diabetes mellitus and bone metabolism. CD38 has been used as a prognostic marker in leukemia.

