

Synonym

Complement C5,C5,CPAMD4

Source

Biotinylated Cynomolgus Complement C5 Protein, His,Avitag(CO5-C82E3) is expressed from human 293 cells (HEK293). It contains AA Gln 19 - Cys 1676 (Accession # [XP_005580972.1](#)).
Predicted N-terminus: Gln 19

Molecular Characterization

Complement C5(Gln 19 - Cys 1676)
XP_005580972.1

Poly-his

Avi

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).
The protein has a calculated MW of 190.1 kDa. The protein migrates as 70 kDa and 120-130 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE).

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated Cynomolgus Complement C5 Protein, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

SEC-MALS




The purity of Biotinylated Cynomolgus Complement C5 Protein, His,Avitag (Cat. No. CO5-C82E3) is more than 85% and the molecular weight of this protein is around 170-210 kDa verified by SEC-MALS. [Report](#)



Biotinylated Cynomolgus Complement C5 Protein, His,Avitag™ (MALS verified)

Catalog # CO5-C82E3



Immobilized Monoclonal Anti-Human C5a Human Antibody, Human IgG1 (Cat. No. CO5-M50) at 2 µg/mL (100 µL/well) can bind Biotinylated Cynomolgus Complement C5 Protein, His,Avitag (Cat. No. CO5-C82E3) with a linear range of 1-63 ng/mL (QC tested).

Background

Derived from proteolytic degradation of complement C5, C5 anaphylatoxin is a mediator of local inflammatory process. C5 precursor is first processed by the removal of 4 basic residues, forming two chains, beta and alpha, linked by a disulfide bond. C5 convertase activates C5 by cleaving the alpha chain, releasing C5a anaphylatoxin and generating C5b (beta chain + alpha' chain). Activation of C5 by a C5 convertase initiates the spontaneous assembly of the late complement components, C5-C9, into the membrane attack complex. C5b has a transient binding site for C6. The C5b-C6 complex is the foundation upon which the lytic complex is assembled. The C5a anaphylatoxin interacts with C5AR1 and tick complement inhibitor. C5a is also a potent chemokine which stimulates the locomotion of polymorphonuclear leukocytes and directs their migration toward sites of inflammation.

