



Synonym

FAP,FAPalpha,SIMP,Seprase,APCE

Source

FITC-Labeled Human FAP, Fc Tag (FAP-HF263) is expressed from human 293 cells (HEK293). It contains AA Leu 26 - Asp 760 (Accession # [Q12884-1](#)). It is the FITC labeled form of Human FAP, Fc Tag (FAP-H5263).
Predicted N-terminus: Pro

Molecular Characterization

Fc(Pro 100 - Lys 330) P01857	FAP(Leu 26 - Asp 760) Q12884-1
---------------------------------	-----------------------------------

This protein carries a human IgG1 Fc tag at the N-terminus.
The protein has a calculated MW of 111.5 kDa. The protein migrates as 115-130 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Conjugate

FITC
Excitation source: 488 nm spectral line, argon-ion laser
Excitation Wavelength: 488 nm
Emission Wavelength: 535 nm

Labeling

The primary amines in the side chains of lysine residues and the N-terminus of the protein are conjugated with FITC using standard chemical labeling method. The residual FITC is removed by molecular sieve treatment during purification process.

Protein Ratio

The FITC to protein molar ratio is 2-4.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please protect from light and avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE




FITC-Labeled Human FAP Protein, Fc Tag

Catalog # FAP-HF263




FITC-Labeled Human FAP, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA



Immobilized Anti-FAP Antibody, Mouse IgG2a (FAP5) at 1 µg/mL (100 µL/well) can bind FITC-Labeled Human FAP, Fc Tag (Cat. No. FAP-HF263) with a linear range of 0.02-0.313 µg/mL (QC tested).

Bioactivity-FACS



2e5 of anti-FAP CAR-293 cells were stained with 100 µL of 3 µg/mL of FITC-Labeled Human FAP, Fc Tag (Cat. No. FAP-HF263) and negative control protein respectively, FITC signal was used to evaluate the binding activity (QC tested).

Background

FAP (also known as seprase) is a Type II transmembrane serine protease. Both plasma membrane and soluble forms exhibit post-proline cleaving endopeptidase activity, with a marked preference for Ala/Ser-Gly-Pro-Ser/Asn/Ala consensus sequences. Degrade also gelatin, heat-denatured type I collagen. Also has dipeptidyl peptidase activity, with a preference for Ala-Pro, Ile-Pro, Gly-Pro, Arg-Pro and Pro-Pro. The plasma membrane form, in association with either DPP4, PLAUR or integrins, is involved in the pericellular proteolysis of the extracellular matrix (ECM), and hence promotes cell adhesion, migration and invasion through the ECM. Promotes glioma cell invasion through the brain parenchyma by degrading the proteoglycan brevican. Acts as a tumor suppressor in melanocytic cells through regulation of cell proliferation and survival in a serine protease activity-independent manner.

Discounts, Gifts,
and more!



FITC-Labeled Human FAP Protein, Fc Tag

Catalog # FAP-HF263

