Catalog # ILC-H5259



#### Synonym

IL-17 RC,IL-17RC,IL17Rhom,IL-17 receptor C,IL-17RL,ZcytoR14

#### Source

Human IL-17 RC, Fc Tag(ILC-H5259) is expressed from human 293 cells (HEK293). It contains AA Leu 21 - His 465 (Accession # <u>NP\_703190.2</u>). Predicted N-terminus: Leu 21

## **Molecular Characterization**

IL-17 RC(Leu 21 - His 465) Fc(Pro 100 - Lys 330) NP\_703190.2 P01857

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 75.9 kDa. The protein migrates as 90-115 kDa when calibrated against <u>Star Ribbon Pre-stained Protein Marker</u> under reducing (R) condition (SDS-PAGE) due to glycosylation.

### Endotoxin

Less than 1.0 EU per  $\mu$ g by the LAL method / rFC method.

# Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

#### Formulation

Lyophilized from 0.22  $\mu$ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

#### Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

#### Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## SDS-PAGE

kDa	M R
180 130 95	Ξ.
65 55	
43	Constant .
33	-
25	-
17	

Human IL-17 RC, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With <u>Star Ribbon Pre-stained Protein Marker</u>).

# SEC-MALS



The purity of Human IL-17 RC, Fc Tag (Cat. No. ILC-H5259) is more than 90% and the molecular weight of this protein is around 200-230 kDa verified by SEC-MALS.



**Bioactivity-ELISA** 

>>> www.acrobiosystems.com





Catalog # ILC-H5259



Immobilized Human IL17A, Tag Free (Cat. No. ILA-H5219) at 2  $\mu$ g/mL (100  $\mu$ L/well) can bind Human IL-17 RC, Fc Tag (Cat. No. ILC-H5259) with a linear range of 5-160 ng/mL (QC tested).



Human IL-17A Protein, Tag Free immobilized on CM5 Chip can Human IL-17 RC, Fc Tag (Cat. No. ILC-H5259) with an affinity constant of 72.8 nM as determined in a SPR assay (Biacore T200) (Routinely tested).

**Bioactivity-BLI** 







Immobilized Human IL-17F, His Tag (Cat. No. ILF-H5244) at 2  $\mu$ g/mL (100  $\mu$ L/well) can bind Human IL-17 RC, Fc Tag (Cat. No. ILC-H5259) with a linear range of 0.02-0.25  $\mu$ g/mL (Routinely tested).



Biotinylated Human IL-17A Protein, His,Avitag<sup>™</sup> (Cat. No. ILA-H82Q1) captured on Biotin CAP - Series S sensor Chip can bind Human IL-17 RC, Fc Tag (Cat. No. ILC-H5259) with an affinity constant of 40.9 nM as determined in a SPR assay (Biacore T200) (Routinely tested).



Loaded Biotinylated Human IL17A, His, Avitag (Cat. No. ILA-H82Q1) on SA Biosensor, can bind uman IL-17 RC, Fc Tag (Cat. No. ILC-H5259) with an

Loaded Human IL-17A&IL-17F Heterodimer Protein, Twin Strep&His Tag (Cat. No. ILF-H52W6) on NTA Biosensor, can bind Human IL-17 RC, Fc Tag





# Human IL-17 RC Protein, Fc Tag (MALS verified)

Catalog # ILC-H5259



affinity constant of 13.8 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

(Cat. No. ILC-H5259) with an affinity constant of 92.3 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Human IL-17F, His Tag (Cat. No. ILF-H5244) on NTA Biosensor, can bind Human IL-17 RC, Fc Tag (Cat. No. ILC-H5259) with an affinity constant of 100 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

## Background

Interleukin-17 receptor C (IL-17 RC), also known as IL17Rhom, IL-17RL, is the receptor for IL17A and IL17F homodimers as part of a heterodimeric complex with IL17RA. IL-17A activity is inhibited by IL-17RA, IL-17F is inhibited by IL-17RC, and a combination of soluble IL-17RA/IL-17RC receptors is required for inhibition of the IL-17F/IL-17A activity. Furthermore, activation of IL17RC can lead to the induction of expression of inflammatory chemokines and cytokines such as CXCL1.



>>> www.acrobiosystems.com

4/21/2025