



Synonym

IL-2 R beta & IL-2 R gamma,IL-2RB & IL-2RG

Source

Mouse IL-2RB&IL-2RG Heterodimer Protein, Fc Tag&Fc Tag (ILG-M5254) is expressed from human 293 cells (HEK293). It contains AA Ala 27 - Glu 240 (IL-2RB) & Trp 23 - Ala 263 (IL-2RG) (Accession # [P16297-1](#) (IL-2RB) & [P34902-1](#) (IL-2RG)).
Predicted N-terminus: Ala 27 (IL-2RB) & Trp 23 (IL-2RG)

Molecular Characterization

IL-2RB (Ala 27 - Glu 240) P16297-1	Fc(Pro 100 - Lys 330) P01857
IL-2RG (Trp 23 – Ala 263) P34902-1	Fc(Pro 100 - Lys 330) P01857

Mouse IL-2RB&IL-2RG Heterodimer Protein, Fc Tag&Fc Tag is produced by co-expression of IL-2RB and IL-2RG, has a calculated MW of 51.5 kDa (IL-2RB) and 54.5 kDa (IL-2RG). Subunit IL-2RB is fused with a human IgG1 Fc tag at the C-terminus and subunit IL-2RG is fused with a human IgG1 Fc tag at the C-terminus. The protein migrates as 65-100 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Mouse IL-2RB&IL-2RG Heterodimer Protein, Fc Tag&Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

SEC-MALS




The purity of Mouse IL-2RB&IL-2RG Heterodimer Protein, Fc Tag&Fc Tag (Cat. No. ILG-M5254) is more than 85% and the molecular weight of this protein is around 138-168 kDa verified by SEC-MALS.
[Report](#)



Mouse IL-2 R beta&IL-2 R gamma Heterodimer Protein, Fc Tag&Fc Tag (MALS verified)

Catalog # ILG-M5254



Immobilized Biotinylated Human IL-2, His,Avitag (Cat. No. IL2-H82E4) at 1 µg/mL (100 µL/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5 µg/well) plate can bind Mouse IL-2RB&IL-2RG Heterodimer Protein, Fc Tag&Fc Tag (Cat. No. ILG-M5254) with a linear range of 0.008-0.25 µg/mL (QC tested).

Background

Both Interleukin-2 receptor subunit beta and Interleukin-2 receptor subunit gamma are receptor for interleukin-2. Common subunit for the receptors for a variety of interleukins. Interacts with SHB upon interleukin stimulation. Probably in association with IL15RA, involved in the stimulation of neutrophil phagocytosis by IL15. This beta subunit is involved in receptor mediated endocytosis and transduces the mitogenic signals of IL2. IL2R exists in 3 different forms: a high affinity dimer, an intermediate affinity monomer (beta subunit), and a low affinity monomer (alpha subunit). The high and intermediate affinity forms also associate with a gamma subunit.

