



Synonym

Integrin alpha 8 beta 1,ITGA8&ITGB1

Source

Human ITGA8&ITGB1 Heterodimer Protein, Twin-Strep Tag&Tag Free(IT1-H5283) is expressed from human 293 cells (HEK293). It contains AA Phe 39 - Leu 1012 & Gln 21 - Asp 728 (Accession # [P53708-1](#) & [P05556-1](#)).  
Predicted N-terminus: Phe 39 & Gln 21

Molecular Characterization

|                                       |          |
|---------------------------------------|----------|
| ITGA8 (Phe 39 – Leu 1012)<br>P53708-1 | Strep II |
| ITGB1 (Gln 21 - Asp 728)<br>P05556-1  |          |

Human ITGA8&ITGB1 Heterodimer Protein, Twin-Strep Tag&Tag Free, produced by co-expression of ITGA8 and ITGB1, has a calculated MW of 116.4 kDa (ITGA8) and 83.7 kDa (ITGB1). Subunit ITGA8 is fused with Strep II-tag at the C-terminus and subunit ITGB1 contains no tag. The protein migrates as 150-180 kDa (ITGA8) and 95-110 kDa (ITGB1) when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under non-reducing (NR) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 150 mM NaCl, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE




Human ITGA8&ITGB1 Heterodimer Protein, Twin-Strep Tag&Tag Free on SDS-PAGE under non-reducing (NR) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

Discounts, Gifts,  
and more!





Immobilized Fibronectin fragment, premium grade (Cat. No. FIN-H5113) at 5 µg/mL (100 µL/well) can bind Human ITGA8&ITGB1 Heterodimer Protein, Twin-Strep Tag&Tag Free (Cat. No. IT1-H5283) with a linear range of 0.01-0.3 µg/mL (QC tested).

Background

Human Integrin alpha 8 beta 1 Heterodimer Protein consists of ITGA8 and ITGB1. The integrin alpha 8 subunit, isolated by low stringency hybridization, is a novel integrin subunit that associates with beta 1. The recently identified alpha 8 integrin subunit associates with beta 1 and is predominantly expressed in smooth muscle and other contractile cells in adult tissues, and in mesenchymal and neural cells during development. In addition, Integrin alpha 8 beta 1 is a receptor for fibronectin and can promote attachment, cell spreading, and neurite outgrowth on fibronectin.

